

Non-Perturbative Renormalization Group Analysis of the Chiral Critical Behaviors in QED

Ken-Ichi Aoki ¹, Keiichi Morikawa, Jun-Ichi Sumi, Haruhiko Terao* ²

and Masashi Tomoyose

Department of Physics, Kanazawa University,
Kakuma-machi, Kanazawa 920-11, Japan

*Dipartimento di Fisica - Università di Genova,
Istituto Nazionale di Fisica Nucleare - sez. di Genova,
Via Dodecaneso, 33 - 16146 Genova, Italy

Abstract

We study the chiral critical behaviors of QED by using Non-Perturbative Renormalization Group (NPRG). Taking account of the non-ladder contributions, our flow equations are free from the gauge parameter (α) dependence. We clarify the chiral phase structure, and calculate the anomalous dimension of $\bar{\psi}\psi$, which is enhanced compared to the ladder approximation. We find that the cutoff scheme dependence of the physical results is very small.

¹e-mail address: aoki@hep.s.kanazawa-u.ac.jp

²On leave of absence from the Department of Physics, Kanazawa University,
e-mail address: terao@dipfis.ge.infn.it

1 Introduction

The dynamical chiral symmetry breaking in gauge theories is one of the most important subjects in particle physics. Many efforts have been devoted to this problem particularly by solving the Schwinger-Dyson (SD) equation for the fermion self-energy mainly in the ladder approximation with the Landau gauge[1]. Adding the four-fermi interactions, the chiral phase diagram and the critical behaviors have been investigated[2] and its characteristic structures with the large anomalous dimension of $\bar{\psi}\psi$ operator has been applied to various models beyond the standard model[3, 4]. In QCD, the ladder SD with the running gauge coupling constant, the improved ladder, gives good results even quantitatively[5]. However the ladder SD is unsatisfactory in many aspects including the strong gauge dependence[6], the difficulty to proceed beyond the ladder[7], no firm base for the improved ladder.

Quite recently, numerical analysis with the Non-Perturbative Renormalization Group (NPRG) [8, 9] draws much attention as a promising new method to study non-perturbative phenomena of the field theory. One of the essential features of the method is that it does not employ any series expansion and therefore is expected to give converging results when improving the approximation, which shows a great contrast against other non-perturbative expansion methods such as $1/N$ or ϵ expansions [10, 11].

Here we briefly explain the general aspects of the NPRG approach. The RG equation is the nonlinear functional differential equation for the Wilsonian effective action, the exact form of which is explicitly written down. To approximate the RG equation, we project it onto a small dimensional subspace of the original full theory space. To improve the approximation, we enlarge the subspace step by step. As the lowest order, we take the local potential approximation (LPA)[12]. The sharp cutoff LPA NPRG has been applied to the dynamical chiral symmetry breaking in gauge theories and it successfully solves the critical behaviors [13, 14]. When we further artificially truncate the LPA down to its ‘ladder parts’, it reproduces the Landau gauge ladder SD results, even the improved ladder results when we adopt the running gauge coupling constant. Thus the LPA NPRG should be regarded as a non-ladder improvement of the ladder SD results, in the course of the systematic approximation of NPRG.

In this article we proceed beyond the LPA analysis of the dynamical chiral symmetry breaking in gauge theories including anomalous dimensions of fields, which brings about two new issues to be carefully dealt with. First, we have to adopt a smooth cutoff scheme to avoid the sharp cutoff singularities for derivative interactions. What we call the cutoff scheme is the profile of a function to suppress the propagation of the low energy modes in the internal lines and is introduced through the momentum dependent mass terms in the

Lagrangian,

$$\frac{1}{2} \int \frac{d^4 q}{(2\pi)^4} \phi(-\mathbf{q}) \cdot \Lambda^2 \cdot C(\Lambda^2/q^2) \cdot \phi(\mathbf{q}), \quad (1)$$

where function C defines the cutoff scheme. In the full space NPRG, physical results do not depend on the cutoff scheme. However the subspace approximation breaks this independence[15], and the cutoff scheme dependence must be checked explicitly.

Second, the NPRG is not manifestly compatible with the gauge symmetries due to the momentum cutoff. Therefore we have to introduce the gauge non-invariant operators to our subspace and constrain them by certain identity, so-called ‘Modified Ward-Takahashi-Slavnov-Taylor Identity’ [16]. This identity ensures the gauge invariance of the total solutions of NPRG.

In the following sections, we formulate the NPRG flow equations beyond the LPA and calculate critical behaviors in QED. Further we investigate the cutoff scheme dependence and the gauge dependence in the covariant α -gauge. These first results beyond the LPA may give us a clue to the total reliability of the NPRG results.

As mentioned above, the ladder SD results have been reproduced in the LPA by further truncating the flow equations. Note that this truncation cannot be regarded as any projection of the RG equation to a subspace. This is crucial for the gauge independence. Therefore the ladder truncation necessarily breaks the gauge invariance. In our system, e.g., the crossed ladder diagrams are included as well as the usual ladder diagrams, keeping the gauge independence of the flows (in some special cutoff scheme), while the ladder SD takes only the ladder parts, and thus suffers from disastrous gauge dependence [6].

2 RG flow equation for the massless QED

We start with the evolution equation for the effective action $\Gamma_\Lambda[A_\mu, \bar{\psi}, \psi]$ [9], where Λ is the infra-red momentum cutoff for the quantum corrections. The effective action is nothing but the one particle irreducible parts of the Wilsonian effective action. The infrared cutoff is introduced by adding the following terms to the bare action:

$$\Delta S_{cut} = \int \frac{d^4 q}{(2\pi)^4} \left\{ \frac{1}{2} A_\mu(-\mathbf{q}) \cdot \Lambda^2 \cdot (\Lambda/q)^{2k} \cdot A_\mu(\mathbf{q}) + (\Lambda/q)^{2k+2} \cdot q_\mu \cdot \bar{\psi}(-\mathbf{q}) \gamma_\mu \psi(\mathbf{q}) \right\}, \quad (2)$$

where a parameter k labels the cutoff scheme and we take the values $k = 1, 2, 3, \dots$. With larger k the cutoff profile becomes sharper and it reaches the θ -function cutoff at $k = \infty$. Since the above cutoff terms maintain the chiral symmetry, the effective action Γ_Λ also respects it. The evolution equation with this cutoff scheme is written as,

$$\Lambda \frac{\partial}{\partial \Lambda} \Gamma_\Lambda = (k+1) \cdot \text{Str} \left\{ \mathbf{C}^{-1} \cdot \left(\mathbf{C}^{-1} + \frac{\overrightarrow{\delta}}{\delta \Phi^T} \Gamma_\Lambda \frac{\overleftarrow{\delta}}{\delta \Phi} \right)^{-1} \right\}, \quad (3)$$

where Φ is a shorthand notation of the fields $\Phi = \{A_\mu, \psi, \bar{\psi}\}$ and $\mathbf{C}^{-1}$ is the matrix,

$$\mathbf{C}^{-1}(\mathbf{q}) \equiv \left(\begin{array}{c|cc} \Lambda^2 \cdot (\Lambda/q)^{2k} \cdot \delta_{\mu\nu} & & \mathbf{0} \\ \hline \mathbf{0} & 0 & (\Lambda/q)^{2k+2} \cdot q_\mu \cdot \gamma_\mu^T \\ & (\Lambda/q)^{2k+2} \cdot q_\mu \cdot \gamma_\mu & 0 \end{array} \right). \quad (4)$$

To approximate the RG equation (3), we project it onto a finite dimensional subspace. Specifying a subspace defines an approximation. Here we take a subspace spanned by the seven chiral invariant operators,

$$\begin{aligned} \Gamma_\Lambda[A_\mu, \bar{\psi}, \psi] = & \int d^4x \left\{ \frac{1}{4} Z_3 F_{\mu\nu}^2 + \frac{1}{2\alpha} (\partial \cdot A)^2 + \frac{1}{2} m^2 \Lambda^2 A_\mu^2 + Z_1 \bar{\psi} \gamma_\mu i \partial_\mu \psi + e A_\mu \bar{\psi} \gamma_\mu \psi \right. \\ & \left. - \frac{1}{2} \frac{G_s}{\Lambda^2} [(\bar{\psi} \psi)^2 - (\bar{\psi} \gamma_5 \psi)^2] - \frac{1}{2} \frac{G_v}{\Lambda^2} [(\bar{\psi} \gamma_\mu \psi)^2 + (\bar{\psi} \gamma_5 \gamma_\mu \psi)^2] \right\}, \end{aligned} \quad (5)$$

where $F_{\mu\nu}$ is the field strength and m^2 is the photon mass which compensates for the lack of the gauge invariance of Γ_Λ .

Note that the chiral and parity invariant 4-fermi interactions without derivatives are spanned by the above two independent operators G_s, G_v . Within the non-derivative (local potential) approximation for multi-fermi operators, the 4-fermi flow equations are not affected at all by other higher dimensional multi-fermi operators, which are composed of at least 8-fermi operators due to the chiral symmetry. In such case, the 4-fermi flows completely determine the critical behaviors for the local potential approximated multi-fermi system, although higher multi-fermi operator flows are necessary when calculating the dynamically generated fermion mass and the chiral condensates [14].

Substituting the effective action (5) to Eq.(3), we obtain the RG flow equations for the dimensionless coupling constants *i.e.* $Z_3, Z_1, \alpha, m^2, e, G_s, G_v$ (see Fig.1).




Figure 1: Diagrams incorporated in our approximation. The diagrams in the solid box correspond to the ladder part and the diagrams in the dashed box are crucial for the gauge invariance.

The effective action (5) actually contains two redundant parameters due to the wave function renormalization. We set Z_1 and Z_3 to be unity by rescaling fields $A_\mu, \psi, \bar{\psi}$. Consequently, the RG equations read,

$$\frac{\partial}{\partial t} m^2 = \left(2 - \frac{e^2}{6\pi^2}\right) m^2 - 4I_1(k)e^2, \quad (6)$$

$$\frac{\partial}{\partial t} e = -\frac{1}{2} \left(3I_4(k, m^2) - I_5(k, m^2, \alpha)\right) e^3 - \frac{1}{12\pi^2} e^3 + 2e (G_S + 4G_V) I_1(k), \quad (7)$$

$$\begin{aligned} \frac{\partial}{\partial t} G_S &= -2G_S - \left(3I_4(k, m^2) - I_5(k, m^2, \alpha)\right) e^2 G_S + 6I_3(k, m^2) e^2 G_S \\ &\quad + 2I_1(k) G_S (3G_S + 8G_V) - 6I_2(k, m^2) e^4, \end{aligned} \quad (8)$$

$$\begin{aligned} \frac{\partial}{\partial t} G_V &= -2G_V - 6I_3(k, m^2) e^2 G_V - \left(3I_4(k, m^2) - I_5(k, m^2, \alpha)\right) e^2 G_V \\ &\quad + I_1(k) G_S^2 - 3I_2(k, m^2) e^4, \end{aligned} \quad (9)$$

$$\frac{\partial}{\partial t} \alpha = \frac{1}{6\pi^2} e^2 \alpha. \quad (10)$$

where t is a cutoff scale parameter defined by $\Lambda = \Lambda_0 e^{-t}$. There appear five independent functions of k, α and m^2 . These functions are defined by the following integrals:

$$I_1(k) = \frac{k+1}{8\pi^2} \int_0^\infty dx x^{2k+2} \left(\frac{1}{1+x^{k+1}} \right)^3, \quad (11)$$

$$\begin{aligned} I_2(k, m^2) &= \frac{k+1}{8\pi^2} \int_0^\infty dx x^{4k+2} \left\{ \left(\frac{1}{1+x^{k+1}} \right)^3 \cdot \left(\frac{1}{1+x^{k+1}+m^2 x^k} \right)^2 \right. \\ &\quad \left. + \left(\frac{1}{1+x^{k+1}} \right)^2 \cdot \left(\frac{1}{1+x^{k+1}+m^2 x^k} \right)^3 \right\}, \end{aligned} \quad (12)$$

$$\begin{aligned} I_3(k, m^2) &= \frac{k+1}{8\pi^2} \int_0^\infty dx x^{3k+2} \left\{ \left(\frac{1}{1+x^{k+1}} \right)^2 \cdot \left(\frac{1}{1+x^{k+1}+m^2 x^k} \right)^2 \right. \\ &\quad \left. + 2 \left(\frac{1}{1+x^{k+1}} \right)^3 \cdot \left(\frac{1}{1+x^{k+1}+m^2 x^k} \right) \right\}, \end{aligned} \quad (13)$$

$$\begin{aligned} I_5(k, m^2, \alpha) &= \frac{k^2-1}{8\pi^2} \int_0^\infty dx \left[x^{2k+1} \left\{ \left(\frac{1}{1+x^{k+1}} \right)^2 \cdot \left(\frac{1}{1+x^{k+1}/\alpha+m^2 x^k} \right) \right. \right. \\ &\quad \left. \left. + \left(\frac{1}{1+x^{k+1}} \right) \cdot \left(\frac{1}{1+x^{k+1}/\alpha+m^2 x^k} \right)^2 \right\} \right. \\ &\quad \left. - x^{3k+2} \left\{ \left(\frac{1}{1+x^{k+1}} \right)^2 \cdot \left(\frac{1}{1+x^{k+1}/\alpha+m^2 x^k} \right)^2 \right. \right. \\ &\quad \left. \left. + 2 \left(\frac{1}{1+x^{k+1}} \right)^3 \cdot \left(\frac{1}{1+x^{k+1}/\alpha+m^2 x^k} \right) \right\} \right], \end{aligned} \quad (14)$$

$$I_4(k, m^2) = \frac{k+1}{k-1} I_5(k, m^2, \alpha=1). \quad (15)$$

Here we note some important features of these RG equations and integrals I_x . The gauge parameter α -dependence of the RG equations comes solely from I_5 , which vanishes with the special cutoff scheme ' $k = 1$ '. Therefore under this cutoff scheme, our RG equations are completely free from α -dependence. Of course this cancellation basically depends on the fact that our beta functions contain a gauge invariant set of diagrams, *e.g.*, the crossed box as well as the box diagrams, and inclusion of the proper anomalous dimension of fields.

The sharp cutoff (large k) limit makes the RG equations singular. The integrals I_1 , I_2 and I_3 approach to finite functions of m^2 , while I_4 and I_5 blow up to infinity, except for the case of vanishing photon mass ($m^2 = 0$). On the other hand, for $m^2 = 0$, integrals $I_4(k, m^2 = 0)$ and $I_5(k, m^2 = 0, \alpha)$ vanish independently of k and α . The photon mass m^2 is to be determined as a function of the gauge coupling constant e by the requirement of the gauge invariance of the effective action $\Gamma_{\Lambda \rightarrow 0}$, and is the order of e^2 . Therefore, the α -dependence of our RG flow equations is the order of e^4 .

We should note the essential difference of our NPRG flows from the perturbative RG. Our RG equations describe the running of the 4-fermi and multi-fermi interactions as well, *e.g.*, $\partial G_S / \partial t \sim G_S^2$, which finally gives the effective potential of the composite operator $\bar{\psi}\psi$ so that we can investigate the dynamical chiral symmetry breaking within our RG equations. Also the beta function of the gauge coupling constant e contains the so-called quadratically divergent diagram contribution, *i.e.*, $\partial e / \partial t \sim e G_S$, which will need special care in relation to the gauge invariance.

3 The critical behaviors of QED

We investigate the $G_S - G_V$ sub-system described by the Eqs.(8) and (9) with the vanishing photon mass approximation ($m^2 = 0$). This system does not suffer from the gauge parameter dependence, and also it has the infinite k limit which completely reproduces the sharp cutoff LPA (Wegner-Houghton equation) results[13, 14]. In more precise treatment with the running photon mass, a small gauge dependence exists, except for the special cutoff profile of $k = 1$. The gauge coupling constant is fixed, no running. Complete analyses with the running gauge coupling constant will be described in the forthcoming paper [17]. Our results here should be compared with the QED SD results with the fixed gauge coupling constant [2]. Then the RG equations determining the critical behaviors are the following coupled differential equations (note $I_3(k, 0) = 1/(8\pi^2)$):

$$\frac{\partial}{\partial t} G_S = -2G_S + 2I_1(k)G_S(3G_S + 8G_V) + \lambda G_S - 6\tilde{I}_2(k, 0)\lambda^2, \quad (16)$$

$$\frac{\partial}{\partial t} G_V = -2G_V - \lambda G_V + I_1(k)G_S^2 - 3\tilde{I}_2(k, 0)\lambda^2, \quad (17)$$

where λ and $\tilde{I}_2$ are defined by $\lambda \equiv 3e^2/4\pi^2$ and $(4\pi^2/3)^2 I_2$ respectively.

In the RG approach, the critical behaviors are governed by the structure of the fixed points. First we consider the case of vanishing gauge coupling constant, where the system is a pure fermi system and we are solving the Nambu-Jona-Lasinio model beyond the ladder (the large N leading) approximation. As is shown in Fig.2, we have two fixed points, the Gaussian infrared fixed point at the origin and the modified Nambu-Jona-Lasinio (NJL) ultraviolet fixed point extended to the vector operator G_V ³. Also there appears another 3rd fixed point at the negative G_S region, which we will ignore here as a fake due to the approximation.




Figure 2: The flow diagram on the $G_S - G_V$ plane with the vanishing gauge coupling constant and with the cutoff scheme ' $k = 1$ '.

There are two phases, the strong phase and the weak phase, and the renormalized trajectory is the straight line defined by $G_S = 8G_V$. In the strong phase, the 4-fermi interactions grow up and diverge at the finite cutoff scale in this truncation. This divergence itself may be regarded as a signal of the limitation of our truncation, and we expect that in an enlarged subspace this singularity will disappear. However, in the local potential approximation, we can still estimate the dynamical fermion mass and the chiral condensate by taking account of the infinite number of higher dimensional fermi operators. In fact, investigating the effective potential of the composite operator $\bar{\psi}\psi$ in either phases, we can conclude that in the strong phase the chiral symmetry is spontaneously broken while in the weak phase the chiral symmetry is respected [14].

Switching on the gauge interactions (Fig.3), these two fixed points move closer to each other and finally meet at some $\lambda (= \lambda_c)$ to pair-annihilate. Beyond this critical gauge coupling constant λ_c , no fixed point exists. The Gaussian fixed point at $\lambda = 0$ moves to

³To be precise, without gauge interactions, there is no ultraviolet fixed point since the 8-fermi operators do not have a fixed point solution.

some finite values of G_S, G_V , which indicates the scale invariant infrared effective 4-fermi interactions generated by the gauge exchange interactions.




Figure 3: The move of the fixed points with the various cutoff schemes. The ladder part LPA results[13, 14] giving the Landau gauge ladder SD equivalents are plotted by the dotted line.

We now understand the total phase structure. For the region $0 \leq \lambda < \lambda_c$, there are two fixed points and two phases divided by the ultraviolet fixed point, the modified NJL fixed point. For $\lambda > \lambda_c$, no fixed point appears and the whole theory space belongs to the strong phase of the dynamical chiral symmetry breaking. These structures generate the peculiar phase diagram on the $\lambda - G_S$ plane. We show the $k = 1$ gauge independent results in Fig.4, where the arrows show the RG flows.




Figure 4: The phase structure on the renormalized trajectory with the cutoff scheme ' $k = 1$ '. We project it onto the $\lambda - G_S$ plane. Note that this is not equal to the section between the critical surface and the $\lambda - G_S$ plane which will be shown in Fig.5

The critical surface obtained in various cutoff schemes are drawn in Fig.5. In Figs.3 and 5, we see that the fixed point position and the critical surface depend largely on the cutoff scheme. This is seen in the RG equations where the criticality is directly determined by the cutoff scheme dependent coefficients (I_1 and I_2). For example, the non-trivial fixed point on the $\lambda = 0$ plane is given by $G_S^* = 8G_V^* = 1/(4I_1(k))$. It should be mentioned that these quantities of the position of the criticality are not the physical quantities at all, that is, they can not be measured. Hence these large cutoff scheme dependences do not cause any difficulty in our approach.




Figure 5: The cutoff scheme dependence of the critical surface in the (G_S, G_V, λ) space. We show the cross sections of the critical surface with $G_S - G_V$ and $\lambda - G_S$ planes. The dotted line is the result of the Landau gauge ladder SD equation.

Our chiral phase criticality qualitatively coincides with the Landau gauge ladder SD results [2]. The quantitative differences come out of two sources. Our results take account of the non-ladder contributions as well as the ladder parts, which guarantees the gauge independence. Also our cutoff profile is smooth while the SD uses the θ -function sharp cutoff. Taking a large k , say $k = 100$, then our results are almost equal to the LPA sharp cutoff results[13, 14] within the line width in every figures.

4 Critical exponent and anomalous dimension

Here, we are going to evaluate the physical quantities, the critical exponent of the ultra-violet fixed point (ν_4), and the anomalous dimension of the fermion mass operator (γ_m). The critical exponent ν_4 is obtained by linearizing the RG flow equations around the ultraviolet fixed point, and it is related to the anomalous dimension of the chiral invariant 4-fermi operators γ_4 through $\nu_4 + 2 = \gamma_4$. On the other hand, the fermion mass operator

$m_f \bar{\psi}\psi$ explicitly breaks the chiral symmetry, and the vanishing mass point is its ultraviolet fixed point. Thus we may evaluate its anomalous dimension γ_m from the linearized RG equation for the operator $m_f \bar{\psi}\psi$,

$$\frac{\partial}{\partial t} m_f = (1 + \gamma_m) m_f + O(m_f^3). \quad (18)$$

The anomalous dimension γ_m is evaluated in terms of the gauge coupling constant λ and the critical scalar 4-fermi interaction $G_S^*(\lambda)$ as

$$\gamma_m(\lambda) = \frac{\lambda}{2} + 8I_1(k)G_S^*(\lambda), \quad (19)$$

where the corresponding diagrams are shown in Fig.6.




Figure 6: Diagrams contributing to the anomalous dimension of the fermion mass operator $\bar{\psi}\psi$.




Figure 7: The critical exponent ν_4 and the anomalous dimension γ_m in various cutoff schemes. The dotted line is the result of the Landau gauge ladder SD equation.

In Fig.7, we plot the critical exponent ν_4 and the anomalous dimension γ_m as a function of the gauge coupling constant λ obtained in various cutoff schemes, with those obtained by the ladder SD equation. Although there are large scheme dependence in the position of criticalities, these exponent and anomalous dimension are almost independent of the cutoff scheme. This indicates that our method and approximation are fairly stable against the cutoff profile dependence.

λ	ν_4			γ_m		
	ladder SD	LPA	our result	ladder SD	LPA	our result
0.0	2.000	2.000	2.000	2.000	2.000	2.000
0.1	1.897	1.918	1.918	1.949	1.987	1.987
0.2	1.789	1.828	1.827	1.894	1.969	1.969
0.3	1.673	1.729	1.726	1.837	1.945	1.944
0.4	1.549	1.619	1.614	1.775	1.914	1.912
0.5	1.414	1.497	1.487	1.707	1.874	1.872
0.6	1.265	1.359	1.342	1.632	1.825	1.820
0.7	1.095	1.200	1.173	1.548	1.763	1.754
0.8	0.894	1.013	0.968	1.447	1.684	1.667
0.9	0.632	0.776	0.699	1.316	1.575	1.542
1.0	0.000	0.418	0.177	1.000	1.396	1.274
at λ_c	0.000	0.000	0.000	1.000	1.170	1.177
(λ_c)	(1.0000)	(1.0409)	(1.0069)	(1.0000)	(1.0409)	(1.0069)

Table 1. The critical exponent ν_4 and the anomalous dimension γ_m . Our gauge independent results ($k = 1$) are compared with the Landau gauge ladder SD[2] and the sharp cutoff LPA[13] results.

Comparing with the ladder SD results, the anomalous dimension γ_m is enhanced much, while the critical exponent ν_4 is almost the same. This large corrections certainly come from the contribution of the non-ladder diagrams. Note that our results are gauge independent due to the inclusion of these non-ladder contributions. In the ladder approximation, these two quantities are not independent and satisfy a relation, $\nu_4 + 2 = 2\gamma_m$, which is given by the simple ladder relation of $\gamma_4 = 2\gamma_m$, where only the scalar 4-fermi operators are taken into account. In our non-ladder extension, this relation is broken and generates the large deviation of γ_m . The numerical values of our gauge independent results ($k = 1$) are listed in Table 1, comparing them with the Landau gauge ladder SD[2] and the sharp cutoff LPA [13] results. It should be noted that the value of the critical gauge coupling constant λ_c depends on the approximation adopted.

5 Discussions and Comments

We solved the next to LPA non-perturbative renormalization group and obtained the critical behaviors of the dynamical chiral symmetry breaking in QED. Our RG system with a special cutoff scheme ($k = 1$) exhibits the gauge parameter independence, which is assured by the inclusion of the non-ladder contributions like the crossed box diagrams.

The NPRG approach to the dynamical chiral symmetry breaking in gauge theories has great potentiality. It turns out that the NPRG with our subspace approximation gives a beyond the ladder approximation restoring the gauge independence. It should be noted that the $k = 1$ cutoff scheme does not work if we go to the higher orders of the derivative expansion, since the beta function integrals do not converge. However, we expect that the gauge dependence should not be large since the NPRG approach may easily incorporate the ‘gauge invariant’ set of diagrams.

We have also confirmed the stability of our method. The cutoff scheme dependences in physical quantities are found to be negligible. Also we get rather small deviations of physical results compared to the LPA. For example, the anomalous dimension γ_m at λ_c is 1.170 in the LPA and 1.177 in our next to LPA respectively. In our level of approximation in this article ($m^2 = 0$), this stability is actually equivalent to the cutoff scheme independence since the infinite k limit corresponds to the sharp cutoff LPA.

Acknowledgment

One of the authors(H. T.) thanks the Department of Physics, University of Genova for hospitality while this paper was completed. K.-I. A. and H. T. are supported in part by Grant-in Aid for Scientific Research (#08240216 and #08640361) from the Ministry of Education, Science and Culture.

References

- [1] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345. T. Maskawa and H. Nakajima, Prog. Theor. Phys. **52** (1974) 1326. R. Fukuda and T. Kugo, Nucl. Phys. **B117** (1976) 250. V. A. Miransky, Nuovo Cim. **90A** (1985) 149.
- [2] K.-I. Kondo, H. Mino and K. Yamawaki, Phys. Rev. **D39** (1989) 2430. K. Yamawaki, in *Proc. Johns Hopkins Workshop on Current Problems in Particle Theory 12, Baltimore, 1988*, eds. G. Domokos and S. Kovesi-Domokos (World Scientific, Singapore, 1988). T. Appelquist, M. Soldate, T. Takeuchi and L.C.R. Wijewardhana, *ibid.* W. A. Bardeen, C. N. Leung and S. T. Love, Phys. Rev. Lett. **56** (1986) 1230. C. N. Leung, S. T. Love and W. A. Bardeen, Nucl. Phys. **B273** (1986) 649.
- [3] K. Yamawaki, M. Bando and K. Matumoto, Phys. Rev. Lett. **56** (1986) 1335. T. Akiba and T. Yanagida, Phys. Lett. **B169** (1986) 432. B. Holdom, Phys. Rev. **D24** (1981) 1441.
- [4] V. A. Miransky, M. Tanabashi and K. Yamawaki, Phys. Lett. **B221** (1989) 177. W. A. Bardeen, C. T. Hill and M. Lindner, Phys. Rev. **D41** (1990) 1647. B. Holdom,

- Phys. Rev. **D54** (1996) 1068. K. Yamawaki, hep-ph/9603293, *in the proceedings of 14th Symposium on Theoretical Physics: Dynamical Symmetry Breaking and Effective Field Theory, Cheju, Korea, 21-26 Jul 1995*.
- [5] K. Higashijima, Phys. Rev. **D29** (1984) 1228. K-I. Aoki, T. Kugo and M. G. Mitchard, Phys. Lett. **B266** (1991) 467. K-I. Aoki, M. Bando, T. Kugo, M. G. Mitchard and H. Nakatani, Prog. Theor. Phys. **84** (1990) 683.
 - [6] K-I. Aoki, M. Bando, T. Kugo, K. Hasebe and H. Nakatani, Prog. Theor. Phys. **81** (1989) 866.
 - [7] K. Kondo and H. Nakatani, Nucl. Phys. **B351** (1991) 236; Prog. Theor. Phys. **88** (1992) 7373. K. Kondo, Int. J. Mod. Phys. **A7** (1992) 7239.
 - [8] K. G. Wilson, I. G. Kogut, Phys. Rep. **12** (1974) 75. F. Wegner, A. Houghton, Phys. Rev. **A8** (1973) 401. J. Polchinski, Nucl. Phys. **B231** (1984) 269. G. Keller, C. Kopper and M. Salmhofer, Helv. Phys. Acta **65** (1992) 32.
 - [9] C. Wetterich, Phys. Lett. **B301** (1993) 90. M. Bonini, M. D'Attanasio, G. Marchesini, Nucl. Phys. **B409** (1993) 441. T. R. Morris, Int. J. Mod. Phys, **A9** (1994) 2411.
 - [10] K-I. Aoki, K. Morikawa, W. Souma, J.-I. Sumi and H. Terao, Prog. Theor. Phys. **95** (1996) 409.
 - [11] M. Alford, Phys. Lett, **B336** (1994) 237. N. Tetradis and C. Wetterich, Nucl. Phys. **B422** (1994) 541. T. R. Morris, Phys. Lett. **B329** (1994) 241.
 - [12] A. Hazenfratz and P. Hazenfratz, Nucl. Phys. **B270** (1986) 269. T. R. Morris, Phys. Lett, **B334** (1994) 355.
 - [13] K-I. Aoki, K. Morikawa, W. Souma, J.-I. Sumi, H. Terao and M. Tomoyose, KANAZAWA-96-21.
 - [14] K-I. Aoki, K. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, KANAZAWA-96-22.
 - [15] R. D. Ball, P. E. Haagensen, J. I. Latorre and E. Moreno, Phys. Lett. **B347** (1995) 80. K-I. Aoki, K. Morikawa, W. Souma, J.-I. Sumi and H. Terao, in preparation.
 - [16] C. Becchi, *On the construction of renormalized quantum field theory using renormalization group techniques*, in: Elementary Particles, Field Theory and Statistical Mechanics, eds. M. Bonini, G. Marchesini, and E. Onofri, Parma University, 1993. U. Ellwanger, Phys. Lett. **B335** (1994) 364. U. Ellwanger, M. Hirsch, and A. Weber,

Z. Phys. **C69** (1996) 687. M. Bonini, M. D’Attanasio, and G. Marchesini, Nucl. Phys. **B418** (1994) 81; **B421** (1994) 429; **B437** (1995) 163; Phys. Lett. **B346** (1995) 87. M. D’Attanasio, and T. R. Morris, Phys. Lett. **B378** (1996) 213. F. Freire, and C. Wetterich, Phys. Lett. **B380** (1996) 337.

[17] K-I. Aoki, K. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, in preparation.